

STAFF SUMMARY SHEET

	TO	ACTION	SIGNATURE (Surname), GRADE AND DATE		TO	ACTION	SIGNATURE (Surname), GRADE AND DATE
1	DFB	sign	Putnam, O6, 30 Sep 14	6			
2	DFER	approve	Solt, AD22, 1 Oct 14	7			
3	DFB	action		8			
4				9			
5				10			

SURNAME OF ACTION OFFICER AND GRADE	SYMBOL	PHONE	TYPIST'S INITIALS	SUSPENSE DATE
Lt Col Maresh	DFB	333-3722	rwm	Not Applicable
SUBJECT Clearance for Material for Public Release				DATE USAFA-DF-PA- 451 20140929

SUMMARY

1. PURPOSE. To provide security and policy review on the document at Tab 1 prior to release to the public.

2. BACKGROUND.

Authors: Hallenbeck PC (LSRC/USAFA), Grogger M (LSRC/USAFA), Mraz M (LSRC/USAFA), Veverka D (LSRC/USAFA).


Title: Building a Better Mousetrap I: using Design of Experiments with unconfounded ions to discover superior media for growth and lipid production by Chlorella sp. EN1234.

Release Information: This manuscript will be submitted for consideration for presentation in the journal Bioresource Technology. DOD disclaimer is located on the last page in Acknowledgements section.

3. DISCUSSION. N/A

4. VIEWS OF OTHERS. N/A

5. RECOMMENDATION. Sign coord block above indicating document is suitable for public release. Suitability is based solely on the document being unclassified, not jeopardizing DoD interests, and accurately portraying official policy.


Lt Col Ryan W. Maresh
Assistant Professor of Biology

Tab
1. Copy of manuscript

Manuscript Number:

Title: Building a Better Mousetrap I: using Design of Experiments with unconfounded ions to discover superior media for growth and lipid production by *Chlorella* sp. EN1234

Article Type: SI: Bio/Chemicals from Algae

Keywords: Biofuels; algae; medium optimization; lipid production; unconfounded ion matrix; Scheffe mix process

Corresponding Author: Prof. Patrick Hallenbeck,

Corresponding Author's Institution:

First Author: Patrick Hallenbeck

Order of Authors: Patrick Hallenbeck; Melanie Grogger; Megan Mraz; Donald Veverka

Abstract: An unconfounded Scheffe Mix approach was used to probe important ions and their interactions in supporting biomass and lipid production by *Chlorella* sp. EN1234. Six major cations and anions; NH_4^+ , NO_3^- , Na^+ , K^+ , PO_4^- and Cl^- ; were investigated. Piepel plots and RSM analysis showed that in a number of cases, the major media anions PO_4^- and Cl^- negatively influence final cell densities, and that maximal cell density is obtained with nitrate over ammonium, with an optimal effect when mixed with equal molar potassium. Although it is commonly assumed that lipid content increases in nitrogen-deficient media, here little correlation was found between nitrogen content and total lipid content with mixtures that supported high lipid productivity. Thus these mixtures define the composition space within which further R&D might produce the best trade-off between total biomass production and high cellular lipid content.

Dear Editor,

We hereby submit to you our manuscript for consideration for publication in Bioresource Technology in the Special Issue: Algal Biofuels and Chemicals. With this letter I would like to attest to the following points:

1. This manuscript should be classed in the section Biofuels from Algae: 30.060
2. That all the authors mutually agree for submitting their manuscript to BITE
3. The manuscript is the original work of the authors,
4. The manuscript has not been submitted earlier to BITE
5. The manuscript is novel in several respects. It describes the use of DOE (design of experiments) to investigate the effects of media components on biomass and lipid production in a way that is very seldom used, an unconfounded ion study. Previous media composition studies have varied ions as their salts, leading to situations where it is impossible to determine if the effect observed is due to the ion of interest, or its counter ion, or simply the ionic strength. This pitfall is avoided in the present study. In addition to determining unique compositions that lead to high levels of biomass and lipid production, we have also investigated several novel aspects. For example, we show that chloride ion is a generalized inhibitor, and how different mixes of sodium and potassium might be useful depending upon the nitrogen source utilized.


I hope that you find our manuscript satisfactory for publication.

Sincerely,


A handwritten signature in black ink, appearing to read 'P. Hallenbeck', with a stylized, cursive script.

Patrick C. Hallenbeck


A



B



C



Highlights:

- An unconfounded mix approach was used to probe important ions & their interactions
- The major media anions PO_4^- and Cl^- negatively influence final cell densities
- With *Chlorella* EN1234, maximal cell density is obtained with nitrate
- Little correlation was found between nitrogen content and total lipid content
- A composition space is defined for the best trade-off in lipid production

**Building a Better Mousetrap I: using Design of Experiments with
unconfounded ions to discover superior media for growth and lipid
production by *Chlorella sp.* EN1234 ^a**

Patrick C. Hallenbeck^{b*}, Melanie Grogger, Megan Mraz, and Donald Veverka,

Life Sciences Research Center, Department of Biology

United States Air Force Academy

2355 Faculty Drive, USAF Academy, Colorado 80840

*Corresponding author:

Email: Patrick.hallenbeck@umontreal.ca

^a*The views expressed in this article are those of the authors and do not reflect the official
policy or position of the United States Air Force, the Department of Defense, or the US
Government.*

^bOn sabbatical leave from the Département de microbiologie, infectiologie, et
immunologie, Université de Montréal, CP 6128, Centre-ville, Montréal. PQ H3C 3J7
Canada

An unconfounded Scheffe Mix approach was used to probe important ions and their interactions in supporting biomass and lipid production by *Chlorella sp.* EN1234. Six major cations and anions; NH_4^+ , NO_3^- , Na^+ , K^+ , PO_4^- and Cl^- ; were investigated. Piepel plots and RSM analysis showed that in a number of cases, the major media anions PO_4^- and Cl^- negatively influence final cell densities, and that maximal cell density is obtained with nitrate over ammonium, with an optimal effect when mixed with equal molar potassium. Although it is commonly assumed that lipid content increases in nitrogen-deficient media, here little correlation was found between nitrogen content and total lipid content with mixtures that supported high lipid productivity. Thus these mixtures define the composition space within which further R&D might produce the best trade-off between total biomass production and high cellular lipid content.

1.0 Introduction

Microalgae are a very diverse group of organisms which display a wide spectrum of potential ability to produce biofuels, nutraceuticals, or other valuable chemicals. In particular, they are under intense study recently for their potential for the sustainable production of biofuels (Abdelaziz et al., 2013 a, b; Fields et al., 2014; Leite & Hallenbeck 2011; Leite et al., 2013; Moody et al., 2014; Lee et al., 2014; li et al., 2014). Variation in microalgal capabilities is linked to the presence and diversity of metabolic pathways. In addition, nutrient composition and ratios might also determine where the cellular metabolism is directed. For example, key nutrients that reportedly affect lipid yield include carbon (in the form of carbon dioxide), nitrogen and phosphorus (Converti et al., 2009; Gordillo et al., 1998; Griffiths et al., 2014 a, b; Sheehan et al., 2008). Algal culture media

are composed of mineral nutrients in the form of salts and a great deal of effort over the past decade has been devoted to finding the optimal concentrations of salts for different species, resulting in a vast number of varying media recipes.

A number of studies have shown that microalgal cellular lipid accumulation, generally in the form of triacylglycerides, is positively affected by stress conditions such as nutrient deficiency or high irradiance (Harwood & Guschina, 2009; Hu et al. 2008; Roessler 1990). Due to the wide diversity of microalgal species and the complexity of cellular biochemical processes involved in carbon flux, it might be expected that there are general mechanisms for regulation of carbon flux that apply to all algae, as well as other, more specific regulatory processes that only apply to specific subgroups. Although many studies have focused on identifying the regulatory mechanisms for lipid accumulation under stress, relatively little is known about the correlation between lipid productivity of microalgae and general growth conditions. Until now, most strains of microalgae have been grown in cultivation media that were developed several decades ago for the general growth of larger groups of algae.

Changes in growth conditions; light intensity (Converti et al., 2009; Xia et al., 2013), temperature (Roleda et al., 2013), pH (Skrupski et al., 2013) and/or nutrient composition (Hu et al., 2008; Leite et al., 2013); can impact both lipid quality and quantity (positively or negatively). For example, a decrease in temperature, while not necessarily affecting the overall lipid yield, changes the lipid composition towards polyunsaturated fatty acids that are less suitable for biodiesel production (Converti et al., 2009). While irradiance can affect lipid production, high irradiance can actually reduce algal growth rate by

1
2
3
4
5 photoinhibition (Radakovits et al., 2010) and thereby reduce lipid productivity overall.
6

7
8 These cellular responses to irradiance and temperature changes are largely species-
9
10 dependent (Guschina et al., 2006).
11
12

13
14 Previously, nutrient manipulations aimed at changing the concentration of a particular ion
15
16 have been achieved mainly by changing the amounts of particular salt constituents of the
17
18 growth medium, neglecting the effect of the corresponding counter ion (Niedz and Evens,
19
20 2007). This leads to a confounding effect where it is not possible to quantitatively separate
21
22 the effects due to the two different ions. As such, the total ion concentration is often
23
24 increased in these studies, creating yet another series of confounded variables. In previous
25
26 work, an experimental design approach was used to investigate the effects of five different
27
28 ions on specific growth rates using a DOE (Design of Experiments) approach with an
29
30 unconfounded ion matrix (Evens and Niedz, 2010). This type of experimental design was
31
32 shown to produce a much richer data set than possible with a one-factor-at-a-time
33
34 approach. The optimization of biofuel production is essential to making renewable fuel
35
36 sources economically viable. Here this approach was used to examine the effects of media
37
38 composition and other environmental manipulations on biomass and lipid accumulation
39
40 with the ultimate goal of formulating a model that identifies the drivers for growth and
41
42 lipid production and predicts the optimal media composition for desired productivity
43
44 indices.
45
46
47
48
49
50
51

52 **2.0 Materials and Methods**

53 *2.1 Strains and growth conditions*

54
55
56
57
58
59
60
61
62
63
64
65

Chlorella sp. EN1234 was obtained from Dr. Juergen Polle, Brooklyn College, and cultures were grown in 3 mL volumes in 12-well covered plates without agitation at 25°C, with a light intensity of 150 μ E on a 16-hour light/8-hour dark cycle. Each 3 mL culture was inoculated with 3,000 algae cells/ μ L.

2.2 Growth, lipid and biomass measurements

Optical density (OD) measurements were taken at 600 nm with a BioTek μ Quant[®] microplate spectrophotometer. Prior to measurement of OD measurement, cultures were agitated on a microtiter plate shaker for 2 minutes to re-suspend any settled algal cells. Variation between runs was \pm 25% and variation between biological duplicates within the same run were \pm 10%. Cell counts and neutral lipid productivity were determined using flow cytometry with a BD Accuri[®] C6 Flow Cytometer. Flow cytometry samples were prepared in 1.5 mL conical bottom tubes with 50 μ L of culture and 150 μ L of deionized water (final sample volume of 200 μ L). Samples were measured without BODIPY[®] 493/503 (Life Technologies, D3922) dye and with 1 μ L of 200X BODIPY[®] (200 ng/ μ L in DMSO). Dyed samples were vortexed and incubated for 10 minutes before measuring. Undyed samples were vortexed and measured immediately.

2.3 Design of Experiments study of operational parameters

Using experimental design software (Design Expert 8) and the mixture and optimal design functions, a matrix was developed to optimize freshwater media. Such mixture designs give response surface designs that are constrained such that all the component proportions add up to 1. Here, a Scheffe Mix Model with a quadratic process order, special cubic mix order and a sixth combined order limit was used. 210 media variations were generated and

used to assess algal biomass and neutral lipid production. Each medium consisted of a basal medium to which were added varying amounts of H_3PO_4 , HCl , HNO_3 , KOH , NaOH , NH_4OH , $\text{Mg}(\text{NO}_3)_2 \times 6\text{H}_2\text{O}$, $\text{MgCl}_2 \times 6\text{H}_2\text{O}$, and $\text{Mg}(\text{OH})_2$ (Table S1).

2.4 Phylogenetic analysis of EN1234

EN1234 is a natural isolate that has been little characterized or described until now. To better understand its relationship to previously described microalgae, portions of its DNA encoding 18S rRNA and 23S rRNA were amplified by PCR and sequenced. Briefly, samples were held at 94°C for 5 minutes; subjected to 30 cycles of 94° for 40 seconds, 61°C for 40 seconds, and 72°C for 90 seconds; and then held at 72°C for 5 minutes before storing at 4°C . PCR reactions were run on an Eppendorf 6325 Mastercycler[®] Pro S PCR machine using an Ambion AgPath-ID One-Step RT-PCR[®] Kit (catalog # AM1005). PCR solution components included: 12.5 μL of 2X RT-PCR Buffer, 1 μL 5X RT-PCR Enzyme Mix, 1 μL of forward primer, 1 μL of reverse primer, 3 μL of algal culture, and 6.5 μL of nuclease-free water (total volume 25 μL). The primers used were: 23S (PRIMER_23SU-1 AGGGGTARAGCACTGYTTYG, PRIMER_23SU-2 CCTTCTCCCGAAGTTACG) and 18S (Forward: GTGGTAACGGGTGACGG, Reverse: GTGCGGCCCAAGACATC). The sequences obtained have been deposited in GenBank (accession numbers: KM213393, 23S, KM213394, 18S). Basic BLAST (NCBI) was used to obtain sequences showing significant homology. These were aligned online using CLUSTALW (<http://www.genome.jp/tools/clustalw/>) and the FASTA output was then used as input to tree construction using the Maximum Likelihood function of MEGA6 (<http://www.megasoftware.net/>).

3.0 Results and Discussion

3.1 Growth of *Chlorella* sp. EN1234 on the 210 solutions

En1234 was originally supplied to us as a potential candidate for lipid production studies. Its phylogenetic relationship to previously studied algae was determined using 18S rRNA and 23S rRNA sequence information obtained as described in Materials and Methods. 18S rRNA sequences showed that it was highly similar to known *Chlorella* species (not shown). The results based on its plastid 23S rRNA sequence, which is more variable than that of the nuclear-encoded 18S rRNA, show that EN1234 is an outlier to known *Chlorella* species, most certainly within the family Oocystaceae and quite possibly a new *Chlorella* (Figure 1). For the purposes of this article it will be called *Chlorella* sp. EN1234.

3.2 Piepel plot analysis of the growth of *Chlorella* sp. EN1234 on the different mixtures

Here the growth of this strain was assessed using the matrix of solutions set up as described in Materials and Methods. The final OD_{600s} obtained for *Chlorella* sp. EN1234 growing on the 210 solutions were fit to a special cubic (mix) linear (process) model. Anova (Table S2) gave a model F-value of 8.18 implying that the model is significant. There is only a 0.01% chance that an F-value this large could occur due to noise. There appeared to be a good fit in terms of other parameters as well, as shown by a normal plot of the residuals and plots of residuals versus predicted and predicted versus actual (Figure S1). The variation in response (OD₆₀₀) expected for changes in single variables while the proportions of the others are held constant was examined using Piepel plots (Figure 1). These are trace plots which allow all the varied components to be viewed at once.

Essentially, the factors tool of the software sets the reference blend through which the traces are plotted, allowing one to visualize how sensitive the formulation is to deviations from the central point (reference blend). Thus, trace plots (also called perturbation plots in response surface and factorial designs) help one to compare the effects of all the components in the design space. In the Piepel plot variation used here, the component under study is varied, while the ratios of the other components are held constant.

When examined using Piepel plots (Figures 2 and 3), the mixtures under study here showed a number of salient points. Firstly, the predicted final cell density (OD_{600}) appeared to be sensitive to total ionic composition with the higher total ions (Fig. 2A) giving a higher OD than the lower ionic compositions (Fig. 2B and 2C). Secondly, as can be seen from the figure, at the lowest ionic concentration (2 mM) there is relatively little variation with changes in any of the parameters (Fig 2C), whereas all the changes are accentuated at 20 mM total ions (Fig. 2A). In all cases, PO_4^- (C) and Cl^- (F) are seen to negatively influence cell density in these mixtures over a relatively small range. With cultures at 20 mM total ions, NH_4^+ and NO_3^- appear to have a negative impact over a longer range (i.e. to higher concentrations).

It was also interesting to examine the Piepel traces obtained with this model with *Chlorella sp.* EN1234 under the conditions that gave the highest response, run 44 (Figure 3). Again, these plots show that the greater the total ion concentration, the greater the final OD_{600} that is obtained (Fig. 3A). These plots suggest that these conditions are optimal since variation of any of the parameters, either to greater or lesser amounts, leads to a drop-off in cell density. In this case, and especially at 20 mM total ions (Fig. 3A), changes in all

of the parameters had the same magnitude effect over most of the range visualized (Except for C and F which do not extend into the negative area since they are already at a minimum).

3.3 RSM plot analysis of the growth of *Chlorella* sp. EN1234 on the different mixtures

While Piepel plots can show roughly how the system as a whole responds, more detailed information about specific interactions between ions can be obtained from surface plots. For this purpose, surface plots under conditions where maximal OD₆₀₀s are obtained were examined for a few ions that provided insights (Figures 4 and 5). When mixtures of ammonium, nitrate, and potassium are varied at 20 mM total ions (Fig. 4A), a number of interesting effects are seen. Maximal cell density is seen with nitrate and little or no ammonium (along the AB axis) but the optimal effect of nitrate is seen when mixed with an equal amount of potassium (Na⁺ constant at 0.312). With no nitrate present (axis AE) very little effect of mixing different proportions of potassium and ammonium is seen, thus this effect is specific to nitrate among the two nitrogen sources. Moreover, ammonium without any nitrate present only poorly supports growth.

Interestingly, when potassium is absent (axis AB) maximum growth is seen with an equal mixture of ammonium and nitrate. On the other hand, very similar analogous effects are seen if potassium is held constant (0.301) and ammonium, nitrate and sodium are varied at 20 mM total ions (Fig. 4B). Thus, as before, nitrate without any ammonium supports maximum cell density, but only in an equal mixture with sodium (BD axis). Again, in analogy with what was found with potassium, in the absence of sodium, an equal

1
2
3
4
5 mixture of ammonium and nitrate is required for maximum cell density (AB axis). Little
6
7 effect of varying mixtures of ammonium and sodium are seen, and in the absence of nitrate
8
9 only poor cell density is seen (AD axis). These effects are only manifested at the high (20
10
11 mM) concentration of ions as identical mixture surfaces at the low ion concentration (2
12
13 mM) show very little changes in contour with variations in proportions (Figure 4C). These
14
15 results suggest that overall charge may be a factor here. Thus, in the absence of one cation,
16
17 equal parts of the other cation and an anion are required. When no anion at all is present,
18
19 things grow poorly, perhaps as much due to the charge difference as to the lack of nitrate
20
21 as a nutrient. These effects are examined further below in section 3.4.
22
23
24
25
26
27

28 Interesting insights can be gained by examining a few other variations in mixture
29
30 components. For example, when ammonium, nitrate and phosphate, the major
31
32 macronutrients for algal growth, were varied at constant sodium and potassium in the
33
34 absence of chloride (Fig 5A), the preference for nitrate over ammonium can be very clearly
35
36 seen (AB axis). While varying the proportions of phosphate and ammonium in the absence
37
38 of nitrate had very little effect on cell density, which remained low (CA axis), changes in
39
40 the relative proportions of phosphate and nitrate had a great effect on biomass which was
41
42 highest when nitrate was highest and phosphate was lowest (CB axis, no ammonium).
43
44 Similarly, it was previously observed that the N:P ratios for optimal growth rates of
45
46 different mixtures are highly dependent upon the ionic context (Evens and Niedz 2010).
47
48
49
50
51
52

53 Holding ammonium constant at zero with intermediate levels of sodium and potassium
54
55 allows the effects of varying proportions of chloride and phosphate on cell density to be
56
57 seen. These had been predicted from the Piepel plots to be important. As indicated in Fig
58
59
60
61
62
63
64
65

1
2
3
4
5
6 5B, chloride and phosphate were both antagonistic with nitrate for biomass production. In
7
8 addition, it can clearly be seen that there were very little interactive effects between
9
10 chloride and phosphate independent of the nitrate concentration. Finally, how the principal
11
12 players, nitrate, sodium, and potassium, interact can clearly be seen in Fig. 5C which
13
14 shows mixtures of these factors at low phosphate (0.01 mM) and the absence of
15
16 ammonium and chloride. Here it can be seen that maximal cell density is obtained at a
17
18 roughly equal mix of these three factors. Changes of this surface with different (lower)
19
20 total ion concentrations suggest that even higher OD₆₀₀s might be achievable with the same
21
22 relative proportions of the varied factors at total ion concentrations above 20 mM.
23
24
25
26
27

28 *3.4 Analysis of lipid production by mixtures showing good growth*

29

30
31 It was of interest to examine the number of mixtures showing good growth, the mixtures
32
33 that led to high cellular lipid contents, and, most importantly, mixtures which led to high
34
35 overall lipid productivity. The number of mixtures supporting good growth decreased
36
37 logarithmically at higher and higher ODs. A final OD₆₀₀ of 0.7 was arbitrarily chosen here
38
39 as providing sufficient growth to be of interest in production. Thirty-seven unique
40
41 mixtures supported growth to this OD or above, twelve of which supported growth to
42
43 OD₆₀₀ ≥ 1.0 (Fig. 6A). Although there is some variation in the composition of the different
44
45 mixtures supporting good growth, in general their characteristics were as determined by
46
47 RSM as described above (Section 3.3). Interestingly, when this criterion is used, this
48
49 *Chlorella* strain does not appear to prefer one nitrogen source over the other as eleven of
50
51 the 37 mixtures contained only NO₃⁻ whereas nine contained only NH₄⁺.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Next, these same mixtures were examined for their effect on cellular lipid content, assessed by FL1 fluorescence as determined by flow cytometry of BODIPY[®] stained cells.

Surprisingly, a very wide variation (over 50 fold for the two extreme cases, mixtures 75 and 106) in lipid content was observed (Table 1 and Fig. 6B). Eight unique mixtures gave a mean FL1 of 50,000 or above (Table 1 and Fig. 6B). Curiously, although it is commonly assumed and often observed that lipid content increases in nitrogen-deficient media, the total nitrogen content of these mixtures varied from 0.55 to 13 mM (Table 1).

Consequently, there appears to be little correlation here between the nitrogen content of the medium and final lipid content. Furthermore, there was no correlation with the N/P ratio as this varied over 400 fold (from 0.15 to 64) in the different mixtures giving high mean FL1 (Table 1). Thus, a low N/P ratio alone cannot explain the high lipid content.

Finally, in terms of potential biodiesel production, total lipid production, i.e. the product of cellular lipid content and total cell numbers, is the most important variable. This was ascertained here by multiplying cellular lipid content (mean FL1 values) by OD₆₀₀, a reasonable proxy for cell numbers. By this measure a number of mixtures gave high lipid productivities (Fig 6C), with the five highest producers (43, 44, 70, 80, 106) not surprisingly found amongst the mixtures giving the highest mean lipid content (Table 1). Accordingly, these mixtures define the composition space within which further R&D might produce the best trade-off between total biomass production and high cellular lipid content.

3.5 Curious cases and oddities.

1
2
3
4
5
6 Finally, the highly varied ionic composition used allows a number of interesting and
7
8 unexpected observations to be made. It should be noted that many of these would not be
9
10 seen with the different media formulations currently in use. Since the method used here
11
12 was explicitly set up to avoid confounding ions, the anions and cations that were varied
13
14 used either H^+ or OH^- as counter ion, and no pH control or buffering was attempted or
15
16 indeed possible. This means that the initial pH of many of the resulting solutions was
17
18 either highly acidic ($56 < pH\ 3$) or highly basic ($26 > pH10$) (Figure S2). However, there
19
20 was a great difference in the behavior at the two different pH extremes. An initial acidic
21
22 pH was very unfavorable for growth with only 5/56 (9%) supporting growth above an
23
24 OD_{600} of 0.2. On the other hand, an initial alkaline pH does not appear to be unfavorable
25
26 since 26/26 (100%) of the mixtures with an initial pH $> pH10$ supported growth above an
27
28 OD_{600} of 0.2, and in fact, 14/26 (54%) had a final OD_{600} above 0.7. This is surprising in
29
30 that *Chlorella* are not noted for being particularly basophilic. Similar effects of different
31
32 initial pHs on growth rates have been previously noted (Evens and Niedz 2010).
33
34
35
36
37
38
39
40

41 The Scheffe Mix Model used here for unconfounded ion mixtures also allowed the
42
43 examination of specific cation and anion requirements, in particular Na^+ , K^+ , and Cl^- ,
44
45 almost universally found in algal media. To our knowledge, freshwater microalgae have
46
47 never been probed for their requirements for these ions. Many media formulations contain
48
49 Cl^- merely because HCl is added to adjust the pH. Here, this was not the case, and 117 of
50
51 the 210 solutions contained no Cl^- . Very surprisingly, this strain grew very well in the
52
53 absence of Cl^- with 23 unique mixtures giving growth to an OD_{600} of 0.8 or better and 9
54
55 unique mixtures supporting growth to an OD_{600} of 1.0 or better (Table 2). In fact, as noted
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5 above (Piepel traces, Figure 2), Cl^- appears to inhibit growth and the highest OD_{600} s were
6
7 obtained in its absence. This is in accordance with a previous study which concluded that
8
9 *C. vulgaris* demonstrates a positive response to cations in its growth medium, but a
10
11 negative response to anions (Evens and Niedz 2010).
12
13
14

15
16 Next, requirements for the important cations Na^+ and K^+ were examined. Of the 113
17
18 mixtures that contained no Na^+ , 19 unique mixtures supported growth to an $\text{OD}_{600} > 0.5$
19
20 with nine unique mixtures supporting growth to an $\text{OD}_{600} > 0.7$ (Table 3). It is noteworthy
21
22 that these nine mixtures all contained no Cl^- , establishing an apparent requirement for the
23
24 absence of this anion in the absence of Na^+ . Of the 105 mixtures where K^+ was absent,
25
26 there were 22 unique mixtures which supported growth to an $\text{OD}_{600} > 0.5$, with ten unique
27
28 mixtures supporting growth to an $\text{OD}_{600} > 0.7$ (Table 3). Similar to what was seen with
29
30 Na^+ , the majority of these mixtures (80%) contained no Cl^- . Not surprisingly, even fewer
31
32 good mixtures were found when the absence of both Na^+ and K^+ cations was probed. There
33
34 were forty-two unique mixtures which lacked both cations, but only 21 of these were even
35
36 capable of supporting growth to an $\text{OD}_{600} > 0.2$, and only 4 could support growth above an
37
38 OD_{600} of 0.4 (Table 3). Interestingly, three of these also lacked Cl^- . Growth of microalgae
39
40 on media simultaneously lacking Na^+ , K^+ , and Cl^- as major ions has not been previously
41
42 observed.
43
44
45
46
47
48
49
50

51 4.0 Conclusions

52
53

54 Here, an unconfounded Scheffe Mix approach and RSM with *Chlorella* sp. EN1234
55
56 showed that the major media anions PO_4^- and Cl^- negatively influence final cell densities.
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 Maximal cell density is obtained with nitrate over ammonium. Several mixtures lacking
7
8 Na^+ , K^+ , and Cl^- as major ions supported significant growth. Little correlation between
9
10 nitrogen content and total lipid content was found with mixtures that supported high lipid
11
12 productivity. Thus, these mixtures define the composition space within which further
13
14 investigation might produce the best trade-off between total biomass production and high
15
16 cellular lipid content.
17
18

19 20 **Acknowledgements** 21

22
23 We thank Dr. T.J. Evens (USDA-Agricultural Research Service, U.S. Horticultural
24
25 Research Laboratory, FL) for initial discussions at the beginning of this project and for his
26
27 help in experimental design, Dr. Juergen Polle (Brooklyn College – CUNY, Brooklyn,
28
29 NY) for providing strain EN 1234 and Dr. Simina Vintila (USAF Academy, Department of
30
31 Biology, Colorado Springs, CO) for technical assistance (making the 210 solutions). PCH
32
33 holds a National Research Council Research Senior Research Associateship Award at the
34
35 Life Sciences Research Center, Department of Biology, United States Air Force Academy.
36
37
38
39
40

41 42 **References** 43

- 44
45 1. Abdelaziz, A.E.M., Leite G.B., Hallenbeck P.C. 2013a. Addressing the challenges for
46
47 sustainable production of algal biofuels: I. Algal strains and nutrient supply,
48
49 Environmental Technology, 34:1783-1805.
50
51
52 2. Abdelaziz, A.E.M., Leite G.B., Hallenbeck P.C. 2013b. Addressing the challenges for
53
54 sustainable production of algal biofuels: II. Harvesting and conversion to biofuels,
55
56 Environmental Technology, 34: 1807-1836.
57
58
59
60
61
62
63
64
65

3. Converti A., Casazza A.A., Ortiz E.Y., Perego P., and Del Borghi M. 2009. Effect of temperature and nitrogen concentration on the growth and lipid content of *Nannochloropsis oculata* and *Chlorella vulgaris* for biodiesel production. Chemical Engineering and Processing, 48:1146-1151.
4. Evens T.J. and Niedz R.P. 2010. Quantification of nutrient-replete growth rates in five-ionhyperspace for *Chlorella vulgaris* (Trebouxiophyceae) and *Peridinium cinctum* (Dinophyceae). European Journal of Phycology 45(3): 247–257.
5. Fields, M.W., Hise, A., Lohman, E.J., Bell, T., Gardner, R.D., Corredor, L., Moll, K., Peyton, B.M., Characklis, G.W., Gerlach, R. 2014. Sources and resources: importance of nutrients, resource allocation, and ecology in microalgal cultivation for lipid accumulation. Applied Microbiology and Biotechnology, 98(11), 4805-4816.
6. Gordillo F.J.L., Goutx M., Figueroa F.L. and Niell F.X. 1998. Effects of light intensity, CO₂ and nitrogen supply on lipid class composition of *Dunaliella viridis*. Journal of Applied Phycology 10:135-144.
7. Griffiths, M.J., van Hille, R.P., Harrison, S.T.L. 2014a. The effect of nitrogen limitation on lipid productivity and cell composition in *Chlorella vulgaris*. Applied Microbiology and Biotechnology, 98(5), 2345-2356.
8. Griffiths, M.J., van Hille, R.P., Harrison, S.T.L. 2014b. The effect of degree and timing of nitrogen limitation on lipid productivity in *Chlorella vulgaris*. Applied Microbiology and Biotechnology, 98(13), 6147-6159.
9. Guschina I.A. and Harwood J.L. 2006. Lipids and lipid metabolism in eukaryotic algae. Progress on Lipid Research 45:160-186.

10. Harwood J.L. and Guschina I.A. 2009. The versatility of algae and their lipid metabolism. *Biochimie* 91(6): 679-684.
11. Hu Q., Sommerfeld M., Jarvis E., Ghirardi M., Posewitz M., Seibert M., and Darzins A. 2008. Microalgal triacylglycerols as feedstocks for biofuel production: Perspectives and advances. *Plant Journal* 54:621–639.
12. Lee, S.-H., Ahn, C.-Y., Jo, B.-H., Lee, S.-A., Park, J.-Y., An, K.-G., Oh, H.-M. 2013. Increased Microalgae Growth and Nutrient Removal Using Balanced N:P Ratio in Wastewater. *Journal of Microbiology and Biotechnology*, 23(1), 92-98.
13. Leite, G.B., Hallenbeck, P.C. Algae Oil, in *Microbial Technologies in Advanced Biofuels Production*, P.C. Hallenbeck, ed. Springer, New York 2011, p. 231-259.
14. Leite, G.B., Abdelaziz A.E.M., Hallenbeck, P.C. Algal biofuels: Challenges and opportunities, *Bioresource Technology*, 145 (2013) 134-141.
15. Li, X., Hu, H.-y., Gan, K., Sun, Y.-x. 2010. Effects of different nitrogen and phosphorus concentrations on the growth, nutrient uptake, and lipid accumulation of a freshwater microalga *Scenedesmus* sp. *Bioresource Technology* 101(14), 5494-5500.
16. Moody, J.W., McGinty, C.M., Quinn, J.C. 2014. Global evaluation of biofuel potential from microalgae. *Proceedings of the National Academy of Sciences of the United States of America*, 111(23), 8691-8696.
17. Niedz R.P. and Evens T.J. 2007. Regulating plant tissue growth by mineral nutrition. In *Vitro Cell Dev Biol – Plant* 43:370-381.

- 1
2
3
4
5
6 18. Radakovits, R., Jinkerson, R.E., Darzins, A., Posewitz, M.C. 2010. Genetic
7
8 Engineering of Algae for Enhanced Biofuel Production. *Eukaryotic Cell*, 9(4), 486-
9
10 501.
11
12
13 19. Roleda, M.Y., Slocombe, S.P., Leahey, R.J.G., Day, J.G., Bell, E.M., Stanley, M.S.
14
15 2013. Effects of temperature and nutrient regimes on biomass and lipid production by
16
17 six oleaginous microalgae in batch culture employing a two-phase cultivation strategy.
18
19 *Bioresource Technology* 129, 439-449.
20
21
22 20. Roessler, P.G. 1990. Environmental control of glycerolipid metabolism in microalgae:
23
24 Commercial implications and future research directions. *Journal of Phycology* 26:393-
25
26 399
27
28
29 21. Skrupski, B., Wilson, K.E., Goff, K.L., Zou, J. 2013. Effect of pH on neutral lipid and
30
31 biomass accumulation in microalgal strains native to the Canadian prairies and the
32
33 Athabasca oil sands. *Journal of Applied Phycology*, 25(4), 937-949.
34
35
36 22. Sheehan J., Dunahay T., Benemann J., and Roessler P.G. 1998. US Department of
37
38 Energy's Office of Fuels Development, July 1998. A Look Back at the US Department
39
40 of Energy's Aquatic Species Program – Biodiesel from Algae, Close Out Report. TP-
41
42 580-24190. Golden, CO: National Renewable Energy Laboratory.
43
44
45 23. Xia, S., Wan, L., Li, A., Sang, M., Zhang, C. 2013. Effects of nutrients and light
46
47 intensity on the growth and biochemical composition of a marine microalga *Odontella*
48
49 *aurita*. *Chinese Journal of Oceanology and Limnology* 31(6), 1163-1173.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure legends

Figure 1. 23S rRNA tree of *Chlorella sp.* EN1234

Distances of *Chlorella sp.* EN1234 to related species were determined as described in Materials and Methods.

Figure 2. Peipel Plot of *Chlorella sp.* EN1234 growing on mixtures containing both NH_4^+ and NO_3^-

Actual component and factor coding was used with Design Expert (DX9) to generate Peipel plots for the mixtures with *Chlorella sp.* EN1234. The components were: A: NH_4^+ , 0.119; B: NO_3^- , 0.119; C: PO_4^- , 0.198; D Na^+ , 0.188; E: K^+ , 0.188; F: Cl^- , 0.188. Traces were done at different total ion concentrations of: A, 20 mM; B, 11 mM; C, 2 mM

Figure 3. Peipel Plot of *Chlorella sp.* EN1234 centered around the best run (Run 44).

Actual component and factor coding was used with Design Expert (DX9) to generate Peipel plots for *Chlorella sp.* EN1234 centered around the best run (highest observed OD_{600}). The components were: A: NH_4^+ , 0.000; B: NO_3^- , 0.377; C: PO_4^- , 0.010; D: Na^+ , 0.312; E: K^+ , 0.301; F: Cl^- , 0.000. Traces were done at different total ion concentrations of: A, 20 mM; B, 11 mM; C, 2 mM

Figure 4. 3D Surface Plots of *Chlorella sp.* EN1234 at constant PO_4^- (0.01 mM)

A. The response of cell density to changes in $[\text{NH}_4^+]$ (A), $[\text{NO}_3^-]$ (B), and $[\text{K}^+]$ (E) at constant Na^+ (0.312), PO_4^- (0.01 mM), and $\text{Cl}^- = 0$. Total ion = 20 mM

B. The response of cell density to changes in $[\text{NH}_4^+]$ (A), $[\text{NO}_3^-]$ (B), and $[\text{Na}^+]$ (C) at constant K^+ (0.301), PO_4^- (0.01 mM), and $\text{Cl}^- = 0$. Total ion = 20 mM

C. The response of cell density to changes in $[\text{NH}_4^+]$ (A), $[\text{NO}_3^-]$ (B), and $[\text{K}^+]$ (C) at constant Na^+ (0.312), PO_4^- (0.01 mM), and $\text{Cl}^- = 0$. Total ion = 2 mM

Figure 5. 3D Surface Plots of *Chlorella* sp. EN1234 with varying PO_4^- or K^+

A. The response of cell density to changes in $[\text{NH}_4^+]$ (A), $[\text{NO}_3^-]$ (B), and $[\text{PO}_4^-]$ (C) at constant Na^+ (0.312), K^+ (0.301), and $\text{Cl}^- = 0$. Total ion = 20 mM

B. The response of cell density to changes in $[\text{NO}_3^-]$ (B), $[\text{PO}_4^-]$ (C), and $[\text{Cl}^-]$ (F) at constant $\text{NH}_4^+ = 0$, Na^+ (0.312), and K^+ (0.251). Total ion = 20 mM

C. The response of cell density to changes in $[\text{NO}_3^-]$ (B), $[\text{Na}^+]$ (D), and $[\text{K}^+]$ (E) at constant $\text{NH}_4^+ = 0$, PO_4^- (0.010), and $\text{Cl}^- = 0$. Total ion = 20 mM

Figure 6. Final OD_{600} s, mean lipid per cell, and total lipid productivity of selected mixtures

The responses of mixtures producing a final OD_{600} of 0.8 or greater are shown. (A) Final OD_{600} , (B) Cellular lipid content as determined by FL1 fluorescence using flow cytometry, (C) total lipid productivity.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Unique Mixtures Giving High Final Lipid Content

Mixture	OD ₆₀₀	Mean FL1	N/P	Total N mM	NH ₄ ⁺ mM	NO ₃ ⁻ mM	PO ₄ ⁻ mM	Na ⁺ mM	K ⁺ mM	Cl ⁻ mM	total ion mM
43	0.91	74100	0.16	1	0.57	0.43	6.33	6.42	0	6.24	20
44	1.18	70800	1.11	0.73	0	0.73	0.66	0	0.61	0	2
59	0.78	71700	5	0.55	0.24	0.31	0.11	3.28	3.64	3.43	11
80	1.56	98300	65.2	13.0	6.32	6.71	0.2	6.15	0.61	0	20
110	0.94	155000	2.12	2.29	0.73	1.56	1.08	2.61	0.98	0.099	7.07
148	0.83	53800	63.0	1.28	0.66	0.62	0.02	0.66	0.039	0.0080	2
200	0.72	76100	0.91	3.42	3.422	0	3.75	3.83	0	0	11
201	0.81	84700	5	1	1	0	0.2	18.8	0	0	20

Table 2 Mixtures with no Cl^- giving good growth

Mixture	pH	OD_{600}	NH_4^+ mM	NO_3^- mM	PO_4^- mM	Na^+ mM	K^+ mM	Cl^- mM	[total ion] mM
78	11.4	1.05	0	0.55	0.11	5.03	5.31	0	11
96	3.73	1.09	0	1.03	0.02	0	0.95	0	2
192	11.25	1.06	3.42	0	3.75	3.83	0	0	11
210	9.46	1.11	0	0.1	0.02	0.94	0.94	0	2
44	10.72	1.18	0	0.73	0.665	0	0.61	0	2
123	7.6	1.20	0.68	0	0.02	0.66	0.63	0	2
207	9.54	1.37	6.50	6.16	0.57	0	6.78	0	20
90	9.49	1.46	6.32	6.71	0.2	6.15	0.61	0	20
4	4.0	1.55	0.073	0.076	0.66	0.59	0.60	0	2

Table 3 Mixtures without Na⁺ and mixtures without K⁺ giving good growth

Mixture	OD ₆₀₀	NH ₄ ⁺ mM	NO ₃ ⁻ mM	PO ₄ ⁻ mM	Na ⁺ mM	K ⁺ mM	Cl ⁻ mM	[total ion] mM
No Na⁺								
44	1.18	0	0.73	0.66	0	0.61	0	2
38	0.873	11.7	0	8.30	0	0	0	20
96	1.13	0	1.03	0.02	0	0.95	0	2
111	0.92	0	5.71	0.11	0	5.18	0	11
141	0.71	0.77	0.0077	0.54	0	14.2	0	15.5
145	0.86	0.67	0.62	0.036	0	0.67	0	2
162	1.47	6.50	6.16	0.57	0	6.78	0	20
186	0.82	0	0.1	0.02	0	1.88	0	2
191	0.79	19.8	0	0.2	0	0	0	20
No K⁺								
175	0.70	1	0	0.2	18.8	0	0	20
202	0.75	0.43	0.12	3.54	3.44	0	3.47	11
199	0.83	3.42	0	3.75	3.83	0	0	11
74	0.86	0.95	0	0.02	1.03	0	0	2
61	0.87	11.7	0	8.30	0	0	0	20
70	0.89	0.57	0.43	6.33	6.42	0	6.24	20
77	0.91	0	5.16	0.11	5.73	0	0	11
120	0.96	5.60	0	0.11	5.29	0	0	11
181	0.98	0	9.82	0.2	9.98	0	0	20
191	0.79	19.8	0	0.2	0	0	0	20
No Na⁺ and K⁺								
204	0.415	1.98	0	0.02	0	0	0	2
169	0.5445	0.1	0	0.02	0	0	1.88	2
191	0.79	19.8	0	0.2	0	0	0	20
38	0.873	11.7	0	8.30	0	0	0	20

Figure

[Click here to download high resolution image](#)






Figure
Click here to download high resolution image

A



B



C




Figure
[Click here to download high resolution image](#)




Figure
Click here to download high resolution image




Figure
[Click here to download high resolution image](#)





Figure
Click here to download high resolution image



Electronic Annex

[Click here to download Electronic Annex: Table S1 Makeup of 210 solutions.xlsx](#)

Electronic Annex

[Click here to download Electronic Annex: Figure S1 Statistics plots EN1234.jpg](#)

Electronic Annex

[Click here to download Electronic Annex: FigS2 pH.tif](#)